Heartsaver® Child CPR Skills Testing Checklist



Student Name Date of Test Scenario: "You are at a park and notice a child suddenly collapse. The scene is safe and you have a cell phone with you but no AED nearby. Demonstrate what you would do next." Assessment and Activation Checks responsiveness Shouts for help/Sends someone to phone 9-1-1 Checks breathing Once student shouts for help, instructor says, "Here's the barrier device. I am going to phone 9-1-1 and get the AED." Cycle 1 of CPR (30:2)				
 *CPR feedback devices preferred for accuracy. Cycle 2 of CPR (repeats steps in Cycle 1) Only check box if step is successfully performed Gives 30 high-quality compressions Gives 2 effective breaths 				
Cycle 3 of CPR (repeats steps in Cycle 1) Gives 30 high-quality compressions	Only check box if step is successfully performed			

Instructor says, "EMS has arrived and is taking over."

STOP TEST

 Instructor Notes Place a check in the box next to each step the student completes successfully. If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation). 					
Test Results Check PASS or NR to indicate pass or needs remediation:					
Instructor Initials	Instructor Number	_ Date			

© 2021 American Heart Association

Heartsaver® Child CPR Skills Testing Critical Skills Descriptors

- 1. Assesses the child and activates emergency response system (this must precede starting compressions) within a maximum of 30 seconds. After determining that the scene is safe:
 - Checks for responsiveness by tapping and shouting
 - Shouts for help and phones 9-1-1
 - · Checks for no breathing or no normal breathing (only gasping)
 - Scans from the head to the chest for a minimum of 5 seconds and no more than 10 seconds
- 2. Cycle 1: Performs high-quality chest compressions (initiates compressions immediately after recognition of cardiac arrest)
 - Correct hand placement
 - Lower half of breastbone
 - 1- or 2-handed (second hand on top of the first) compressions
 - Compression rate of 100 to 120/min
 - Delivers 30 compressions in 15 to 18 seconds
 - Compression depth and recoil—compress at least one third the depth of the chest, about 2 inches (5 cm)
 - Use of a commercial feedback device/manikin is highly recommended
 - Complete chest recoil after each compression

3. Cycle 1: Provides 2 breaths by using a barrier device

- Opens airway adequately
 - Uses a head tilt-chin lift maneuver
- Delivers each breath over 1 second
- Delivers breaths that produce visible chest rise
- Avoids excessive ventilation
- · Resumes chest compressions in less than 10 seconds
- 4. Cycle 2: Performs same steps for compressions and breaths as in Cycle 1
- 5. Cycle 3: Performs same steps for compressions and breaths as in Cycle 1